

Digestive Specialists, Inc. Digestive Endoscopy Center, LLC

www.digestivespecialists.com

Wireless Capsule Endoscopy Prep Instructions

Please read these instructions thoroughly to ensure a successful preparation for your procedure.

If you have the following you cannot have the procedure:

- Heart Loop monitor
- Defibrillator
- Spinal Cord Stimulator

Please let your doctor know if you:

- Ever had a Small Bowel Blockage
- Any Difficulty Swallowing
- Have a **Slow Emptying** stomach
- Take Iron

Capsul	e Prep
--------	--------

- 1. Milk of Magnesia
 - 1 bottle

Date	Time of Arrival
Sugarcreek Twsp	4340 Clyo Road, Suite 200
□ Dayton	1530 Needmore Road, Suite 101
Huber Heights	5697 Shull Road
Springboro	77 W. Eleanor Drive
If you have questions, call the Medical Assistant, at	
After 5:00 PM or on weel	kends, call (937) 534-7330 to page your doctor.

Prep Instructions - Please read line by line to ensure a successful preparation

Day before your capsule procedure – Date: _____

- You may have a normal breakfast and lunch.
 No solid food after 1:00 pm.
- Take only clear liquids after 1:00 PM. (See Clear Liquids Diet on the right.) At a minimum drink fluids as outlined below:

Time	Instructions
5:00 PM	Drink a minimum of 8 oz of clear liquid
6:00 PM	Drink a minimum of 8 oz of clear liquid
7:00 рм	Drink a minimum of 8 oz of clear liquid and take 4 Tablespoons of Milk of Magnesia

Clear Liquids Diet

- Gatorade is preferred
- Clear fruit juices, white grape juice and apple juice
- Kool-Aid or PowerAde
- Clear soup, broth or bouillon
- Water
- Tea or coffee **No cream**
- Hard candies
- Soda pop
- Jell-O or popsicles
- Nothing red or purple
- Nothing alcoholic

8:00 PM Drink a minimum of 8 oz of clear liquid 9:00 PM Drink a minimum of 8 oz of clear liquid

- You may continue on your clear liquid diet until 12:00 MIDNIGHT then nothing until instructed.
- If you are on diabetic medications, you will need specific instructions. Call your doctor's medical assistant if you did not receive any instructions.
- 1 hour before you are scheduled to have your procedure, take the following with a small amount of water only if a box is checked:

☐ 1 Reglan tablet☐ 1 Erythromycin tablet

- Do not take any medications the morning of the test, except the tablet of Reglan or Erythromycin, if ordered. You may take your morning medications four hours after you swallow the capsule.
- No iron supplements five days prior to the capsule endoscopy.

General Information

- Wear loose fitting clothing that is 2 pieces.
- You will need to return to the same office at the end of the day to return the belt.
- It takes approximately 1-2 weeks for the results of the test.
- After the belt is removed you may resume normal activities, to include eating.
- Any questions after the procedure is completed should be directed to your ordering doctor or their medical assistant.