

Digestive Specialists, Inc. Digestive Endoscopy Center, LLC

www.digestivespecialists.com

2 Day Miralax Prep Instructions

Please read these instructions thoroughly to ensure a successful preparation for your procedure.

Date	Time of Arrival	Colonoscopy Prep			
Sugarcreek Twp.	4340 Clyo Road, Suite 100	MiraLAX 238 gm - 1 bottle Dulcolax (bisacodyl)			
☐ Dayton	1530 Needmore Road, Suite 100				
Huber Heights	5697 Shull Road	Laxative			
☐ Springboro	77 W. Eleanor Drive	- 2 tablets			
Miami Valley Hospita	One Wyoming Street • Dayton, 45409	☐ 1 bottle of Magnesium			
☐ Kettering Medical Center 3535 Southern Boulevard • Kettering, 45429 Citrate (only if checked)					
☐ Sycamore Medical Center 4000 Miamisburg Centerville Rd. Miamisburg, 45342					
You are scheduled with Dr					
The Medical Assistant may be reached at					
After 5:00 PM or on weekends, call (937) 534-7330 to page your doctor.					

Important Information

- For 7 days prior to your procedure stop taking iron supplements.
- For 3 days prior to your procedure drink plenty of fluids to hydrate yourself
- Take heart, blood pressure or seizure medications with a small amount of water up to 2 hours before your procedure.
- Complete the entire preparation per the instructions. If you get nauseated or vomit when drinking the prep and you have not finished the entire amount:
 - Stop drinking any further prep or liquids for the next hour until nausea/vomiting has subsided then restart drinking it until finished.
 - If you still cannot finish the prep, please call us at the number listed above.
- Bring a licensed driver. You will not be allowed to drive home.
- You may not take a cab alone.
- Do not smoke on the day of your procedure.



On	Orink 1 bottle of Magnesium Citrate (over the c	ys prior – only if box is checked) counter) at 6 pm	
On	(Day before	ore your procedure)	
	Begin clear liquid diet as instructed on the right. Do not eat any solid foods until after your procedure is completed Drink as much clear liquid as you can all day in order to remain well hydrated. Mix one bottle of Miralax (238gm) in 3 ½ quarts (112 oz) of Gatorade (not red or purple) to make your prep solution At 12:00 Noon, take 2 Dulcolax tablets Between 4:00 PM-6:00 PM drink seven 8 oz glasses of the Miralax prep solution. Shake well before pouring		Clear Liquids Diet (Liquids you can see through) Gatorade(is preferred) Clear fruit juices, white grape juice and apple juice Kool-Aid or PowerAde Clear soup, broth or bouillon Water Tea or coffee - No cream (May have sugar) Hard candies Soda pop Jell-O or popsicles Nothing red or purple No alcoholic beverages No protein drinks or smoothies
2 nd Ha	alf of Miralax Prep - Even if you are po	assing clear liquid the night bo	efore, you still need to complete
the pro	<i>ep.</i> At	ам / рм (6 hours be	fore your arrival time)
•	Drink another seven 8oz glasses of the Nover the next 2 hours.	Miralax prep solution	

After completing the second part of the prep: Do not drink anything else. Do not chew gum or eat

hard candy.