

## Digestive Specialists, Inc. Digestive Endoscopy Center, LLC

www.digestivespecialists.com

## **2 Day OCL Prep Instructions**

**Colonoscopy Prep** 

Please read these instructions thoroughly to ensure a successful preparation for your procedure.

| Date  | Time of Arrival               | <ol> <li>OCL Solution         <ul> <li>prescription required</li> </ul> </li> </ol> |  |  |
|---|-------------------------------|---|--|--|
| Sugarcreek Twp.   | 4340 Clyo Road, Suite 100     | 2. Dulcolax (bisacodyl)   |  |  |
| □ Dayton  | 1530 Needmore Road, Suite 100 | Laxative  – 2 tablets   |  |  |
| ☐ Huber Heights 5697 Shull Road   |                               | ☐1 bottle of Magnesium  |  |  |
| ☐ Springboro  | 77 W. Eleanor Drive           | Citrate (only if checked)   |  |  |
| ☐ Miami Valley Hospital One Wyoming Street • Dayton, 45409                  |                               |   |  |  |
| ☐ Kettering Medical Center 3535 Southern Boulevard • Kettering, 45429       |                               |   |  |  |
| ☐ Sycamore Medical Center 4000 Miamisburg Centerville Rd. Miamisburg, 45342 |                               |   |  |  |
|   |                               |   |  |  |
| You are scheduled with Dr   |                               |   |  |  |
| The Medical Assistant may be reached at                                     |                               |   |  |  |
| After 5:00 PM or on weekends, call (937) 534-7330 to page your doctor.      |                               |   |  |  |
|   |                               |   |  |  |

## **Important Information**

- For 7 days prior to your procedure stop taking iron supplements.
- For 3 days prior to your procedure drink plenty of fluids to hydrate yourself.
- Take heart, blood pressure or seizure medications with a small amount of water up to 2 hours before your procedure.
- Complete the entire preparation per the instructions on the back. If you get nauseated or vomit when drinking the prep and you have not finished the entire amount:
  - Stop drinking any further prep or liquids for the next hour until nausea/vomiting has subsided then restart drinking it until finished.
  - If you still cannot finish the prep, please call us at the number listed above.
- Bring a licensed driver. You will not be allowed to drive home.
- You may not take a cab alone.



• Do not smoke on the day of your procedure.

| On                | Take 1 Magnesium Citrate (Over the cou   | orior, only if box checked) Inter) at 6 pm.   |
|-------------------|--|---|
| On                | (day before  | e your procedure  |
| 2 <sup>nd</sup> H | Begin clear liquid diet as instructed on the right.  Do not eat any solid foods until after your procedure is completed  Drink as much clear liquid as you can all day in order to remain well hydrated.  At 12:00 Noon, take 2 Dulcolax tablets  At 4:00PM mix the prep solution by adding water to the fill line. If you prefer to drink it cold, please mix and refrigerate up to 4 hours prior to drinking.  Between 4:00- 6:00 PM drink eight 8 oz glasses of the prep solution | Clear Liquids Diet (Liquids you can see through)  • Gatorade is preferred • Clear fruit juices, white grape juice and apple juice • Kool-Aid or PowerAde • Clear soup, broth or bouillon • Water • Tea or coffee - No cream (May have sugar) • Hard candies • Soda pop • Jell-O or popsicles • Nothing red or purple • Nothing alcoholic • No protein drinks or smoothies |
| prep.             | and a court of even if you are passing   | areaquia anegire sejore, you am need to complete the  |
| On _              | at   | AM / PM (6 hours before your arrival time)  |

After completing the second part of the prep: Do not drink anything. Do not chew gum or eat hard candy.



Drink another eight 8 oz glasses of the prep solution over the next two hours.